

The Some Lygaeidae (Hemiptera: Heteroptera) Family Species by Collected by Light Traps in Western Turkey

İnanç ÖZGEN¹

Semih ÖRGEL²

Ayşenur TAN³

¹ Fırat University, Engineering Faculty, Bioengineering Department, Elazığ, TURKEY

² Celal Bayar University, Demirci Vocational School, TR-45600, Demirci, Manisa, TURKEY

³ Sakarya University of Applied Sciences, Pamukova Vocational School, Department of Landscape and Ornamental Plants, Sakarya, TURKEY

Corresponding author E-mail: inancozgen@gmail.com

ABSTRACT: This study was carried out between 2013 to 2017 years in Western Turkey. The species were collected by light traps. Totally; The seven species were recorded in Lygaeidae Family. These are: *Remaudierea annulipes* (Barensprung, 1859), *Eremocoris abietis* (L., 1758), *Beosus quadripunctatus* (Müller, 1766), *Nysius cymoides* (Spinola, 1837), *Lamprodema maura* (Fab., 1803), *Metopoplax origani* (Kolenati, 1845) *Camptocera glaberrima* (Walker, 1872). All of specimens were faunistically important for their fauna.

KEYWORDS: Lygaeidae, Western Turkey, New Faunistic Records.

To cite this article: Özgen, İ., Örgel, S., Tan, A., 2021, The Some Lygaeidae (Hemiptera: Heteroptera) Family Species by Collected by Light Traps in Western Turkey, *J..Het.Turk.*, 3 (1):81-83

DOI:10.5281/zenodo.4841114

To link to this article: <https://www.j-ht.org/wp-content/uploads/2021/05/V31-A9.pdf>

Received: Apr 27, 2021; **Revised:** May 22, 2021; **Accepted:** May 27, 2021; **Published online:** May 31, 2021

INTRODUCTION

The Lygaeidae are a family in the Hemiptera (true bugs), with more than 110 genera in 4 subfamilies. The family includes the insects commonly known as milkweed bugs, and also some of those known as seed bugs (Dellapé et al., 2019). Although there have been local studies in Western Turkey, there has not been any detailed study on this area up to now. The studies of Aysev (1974), Lodos et al. (1978) and

Çakır & Önder (1990), Abacıgil et al., (2010); are the other important studies on fauna of Turkish Lygaeidae. During this time; Totally 149 species were found in 59 genera and in 12 families in Central Anatolia and Western Black Sea Regions locations (Lodos et al. 1999). In the last time study on Lygaeoidea; three species were recorded from Turkey: *Emblethis solitarius* Jakovlev, 1882; *Orsillodes longirostris* Puton, 1884 (both Rhyparochromidae) and *Horvathiolus*

kiritschenkoi kiritschenkoi Josifov, 1965 (Çerçi and Tezcan, 2020). Studies on this family need to be detailed in Western part of Turkey.

MATERIAL AND METHODS

The Lygaeidae species were collected in different habitats in İzmir, Aydın, Denizli and Manisa provinces. One light trap on the ground was used for each area from the mid June to the mid September. A 20 watt Philips energy saver white day light bulb was used at each trap and traps were cleared at two weeks intervals. The specimens were identified by Armand Matocq (Paris/France).

RESULTS

Remaudiereana annulipes (Barensprung, 1859)

Material examined: Manisa, Alaşehir, 01.08.2013, 2 male, 1 female, **Totally:** 3 exc.

Distribution in Turkey: Adana, Balıkesir, İzmir, İstanbul (Abacıgil et al., 2010; Yazıcı et al., 2015; Anonymous, 2021).

Note: New record in Manisa province.

Eremocoris abietis (Linnaeus, 1758)

Material examined: İzmir, Seferihisar, 15.08.2013, 4 male, 1 female, **Totally:** 5 exc.

Distribution in Turkey: Bitlis (Tatvan) (Pericart, 1998).

Note: Second record in Turkey fauna.

Beosus quadripunctatus (Müller, 1766)

Material examined: Manisa, Alaşehir, 15.07.2013, 3 male, **Totally:** 3 exc.

Distribution in Turkey: Widespread (Önder et al., 2006; Matocq & Özgen, 2010; Özgen et al., 2018).

Nysius cymoides (Spinola, 1837)

Material examined: Manisa, Alaşehir, 01.08.2013, 12 male, 13 female, **Totally:** 25 exc

Distribution in Turkey: Widespread (Özgen et al., 2020).

Lamprodema maura (Fabricius, 1803)

Material examined: Manisa, Alaşehir, 01.07.2013, 15 male, 6 female, **Totally:** 21 exc

Distribution in Turkey: Balıkesir, Diyarbakır, Edirne (Abacıgil et al., 2010; Tezcan et al., 2010).

Metopoplax origani (Kolenati, 1845)

Material examined: İzmir, Seferihisar, 15.08.2013, 5 male, **Totally:** 5 exc

Distribution in Turkey: Widespread (Çerçi et al., 2018).

Camptocera glaberrima (Walker, 1872)

Material examined: İzmir, Seferihisar, 15.05.2013, 14 male, 7 female, **Totally:** 21 exc

Distribution in Turkey: Adana, Diyarbakır, Kahramanmaraş, Hatay, Gaziantep, Siirt (Önder & Adıgüzel, 1979; Çağatay, 1985; Önder et al., 2006; Matocq & Özgen, 2010).

ACKNOWLEDGEMENTS

The authors are thankful to expert for identification of the Lygaeidae species to Armand Matocq (Paris/France).

REFERENCES

- Abacıgil, T. Ö., Varlı, S. V. ve S. Tezcan., 2010, Edremit (Balıkesir) Körfezi çevresindeki zeytin bahçelerinde kışlak tuzaklarla saptanan Heteroptera türleri. *Türkiye Entomoloji Dergisi*, 34 (1): 105-115.
- Anonymous, 2021, <http://dogalhayat.org/property/lygaeidae-remaudiereana-annulipes/>. Date: 15.04.2021.
- Aysev, N., 1974, *Ege Bölgesi Lygaeidae Familyası Üzerinde Sistematik Çalışmalar*. Güven Matbaası, Ankara, 149 pp.
- Çağatay, N., 1985, Türkiye Rhyparochrominae (Heteroptera-Lygaeidae) alt familyasının taksonomisi ve erkek genital organının önemi üzerinde çalışmalar [Studies on the taxonomy and the morphology of male genitalia of Rhyparochrominae (Heteroptera-Lygaeidae) from Turkey. *Bitki Koruma Bülteni*, 5 (3-4): 73-92.
- Çakır, S., Önder, F., 1990, Türkiye Geocorinae (Het.: Lygaeidae) altfamilyası üzerinde

- sistematik ve faunistik çalışmalar. *Türk. Entomol. Derg.*, 14: 37-52.
- Çerçi, B., Özgen, İ., Dioli, P., 2018, Additional Faunistic Notes on Heteroptera (Hemiptera: Insecta) in East Anatolia (Turkey), *Journal of Entomology and Zoology Studies*, 2018; 6(1): 1225-1231.
- Çerçi, B., Tezcan, S., 2020, Contribution to the Lygaeoidea (Heteroptera) fauna of Turkey based on specimens preserved in the Lodos Entomological Museum, Turkey (LEMT) and additional information. *Journal of Insect Biodiversity*, 15(1): 6-15.
- Dellapé, Pablo M., Henry, Thomas J., 2019, "Family Lygaeidae Schilling, 1829". Lygaeoidea Species File. Retrieved 2019-06-19.
- Yazıcı, G., Yıldırım, E., Moulet, P., 2015, Contribution to the knowledge of the Lygaeoidea (Hemiptera, Heteroptera) fauna of Turkey, *Linzer biol. Beitr.*, 47/1: 969-990.
- Lodos, N., Önder, F., Pehlivan, E., Atalay, R., 1978, *Ege ve Marmara Bölgelerinin Zararlı Böcek Faunasının Tespiti Üzerinde Çalışmalar [(Curculionidae, Scarabaeidae (Coleoptera); Pentatomidae, Lygaeidae, Miridae (Heteroptera)]*. T.C. Gıda, Tarım ve Hayvancılık Bakanlığı, Zir. Müc. Zir. Kar. Gen. Md. Yay., Ankara, 301 pp.
- Lodos, N., Önder, F., Pehlivan, E., Atalay, R., Erkin, E., Karsavuran, Y., Tezcan, S., Aksoy, S., 1999, *Faunistic Studies on Lygaeidae (Heteroptera) of Western Black Sea, Central Anatolia and Mediterranean Regions of Turkey*. Ege Üniversitesi Yayınları, İzmir, 58 s.
- Matocq, A., Özgen, İ., 2010, A Preliminary List of Heteroptera Collected in Mardin and Siirt Provinces from South-Eastern Anatolia of Turkey (Hemiptera). *Mun. Ent. Zool.*, 5 (suppl.): 1011-1019.
- Péricart, J., 1998, *Hémiptères Lygaeidae euro-méditerranéens, volume 2. Faune de France*. Vol. 84B. Fédération Française des Sociétés de Sciences Naturelles, Paris, 453 pp.
- Tezcan, S., Gülperçin, N., Fent, M., 2010, Contribution to the knowledge of the light trap collected Heteroptera fauna occurring in cherry orchards in western Turkey. *Linzer biol. Beitr.* 42/1: 817-823.
- Önder, F., Adigüzel, N., 1979, Some Heteroptera collected by light trap in Diyarbakir (Turkey). *Türkiye Bitki Koruma Dergisi*, 3 (1): 25-34.
- Önder, F., Karsavuran, Y., Tezcan, S., Fent, M., 2006, *Türkiye Heteroptera (Insecta) Kataloğu [Turkey Heteroptera (Insecta) catalogue]*. Meta Basım Matbaacılık Hizmetleri, Bornova-Izmir, 164 p.
- Özgen, İ., Çerçi, B., Kaya, C., 2018, Heteroptera (Hemiptera) Species Determined in Pistachio Orchards of Siirt Province with a New Record for Fauna of Turkey: *Yotvata nergal* Linnavuori, 1993. *Cercetări Agronomice în Moldova*. Vol. LI, No. 4 (176) / 2018: 87-95.
- Özgen, İ., Kara, B.K., Miroğlu, S., Dioli, P., 2020, A New Potential Pest of East and South-Eastern Anatolia in Turkey: *Nysius cymoides* (Spinola, 1837) (Heteroptera, Lygaeidae). *Mun. Ent. Zool.* Vol. 15, 1: (265-267)